**Course Code : BSCC2001** 

**Course Name: Organic Chemistry I** 

# B&EYER STRAIN THEORY

GALGOTIAS UNIVERSITY

Name of the Faculty: Dr. Diwakar Chauhan

**Course Code : BSCC2001** 

**Course Name: Organic Chemistry I** 

### **Baeyer strain theory**

- A The bond angel is of 109° 28' (or 109.5°) for carbon atom in tetrahedral geometry (methane molecule).
- Rever observed different bond angles for different cycloalkanes and also observed some different properties and stability.
- On this basis, he proposed angle strain theory.

Course Code : BSCC2001

**Course Name: Organic Chemistry I** 

### **Conformations of Cycloalkanes**

Rever, when the carbon atoms form part of a ring, the angle will be controlled by the requirements of the ring.

Research BOND ANGLE=

(n – 2) × 180°

[where **n** is no of carbon in ring]



**Course Name: Organic Chemistry I** 


**BOND ANGLES** in some cycloalkanes are as follows: 60 degree 90 degree 108 degree 120 degree IVERSITY

**Course Code : BSCC2001** 

**Course Name: Organic Chemistry I** 

### **Stability of Cycloalkanes**

- Bayer (1885) postulated a theory of angle strain for cycloalkanes in which the difference between a tetrahedral angle (109.5 °) and an internal angle of the appropriate polygon is used as a measure of stability.
- The deviation of bond angles from the tetrahedral angle causes the molecule to be strained and hence unstable compared with molecules with tetrahedral bond angles



**Course Code : BSCC2001** 

**Course Name: Organic Chemistry I** 

ANGLE STRAIN is Angle through which each of valency bond was deflected from normal direction in formation of rings. This is called as angle strain which determine stability of the ring

Angle strain = 1/2 (109°28' – x) where x= internal bond angle GALGGOTIAS UNIVERSITY

**Course Code : BSCC2001** 

**Course Name: Organic Chemistry I** 



#### Table 1.2 : Angle Strain In CycloAlkanes

| Compound     | Bond Angle       | Angle Strain       |
|--------------|------------------|--------------------|
| Cyclopropane | 60 <sup>°</sup>  | 24 <sup>°</sup> 44 |
| Cyclobutane  | 90 <sup>°</sup>  | 9°44               |
| Cyclopentane | 108°             | 0°44               |
| Cyclohexane  | 120 <sup>0</sup> | -5°16              |



**Course Code : BSCC2001** 

**Course Name: Organic Chemistry I** 

### ST&BILITY AND ANGLE STRAIN RELATION

(Stability)

Cyclopropane Cyclobutane Cyclopentane and goes on


UN (Angle strain)

Name of the Faculty: Dr. Diwakar Chauhan

Course Code : BSCC2001

**Course Name: Organic Chemistry I** 

Cyclopropane reaction



Name of the Faculty: Dr. Diwakar Chauhan

**Course Code : BSCC2001** 

**Course Name: Organic Chemistry I** 

## TYPES OF STRAINS

The stability of cycloalkanes are influenced by a combination of three factors:


RAngle strain Torsional strain Steric strain

Course Code : BSCC2001

**Course Name: Organic Chemistry I** 

### Angle strains

Angle through which each of valency bond was deflected from normal direction in formation of rings . This is called as angle strain which determine stability of the ring




Name of the Faculty: Dr. Diwakar Chauhan

**Course Code : BSCC2001** 

**Course Name: Organic Chemistry I** 

### **Torsional strain**



Name of the Faculty: Dr. Diwakar Chauhan


**Course Code : BSCC2001** 

**Course Name: Organic Chemistry I** 


### STERIC STRAIN

The strain due to the repulsive interactions when atoms approach each other too closely. There is increase in energy when atoms are forced too close to one another

 Steric strain focuses on the strain of functional groups bound due to size onto the cyclic ring.
Because rings don't rotate freely in space like linear chains then energy is higher when they are on the same side. For example: cyclopropane



**Higher Energy** 



Lower Energy

#### **Course Code : BSCC2001**

### Why does Baeyer theory fails?

#### **R** The Baeyer theory fails for larger rings because:

- •The angles that Baeyer used for each ring were based on the assumption that the rings of all cycloalkanes are planar.
- In fact, all cycloalkanes except cyclopropane are not planar (flat).

•The reality is that cycloalkanes tend to adopt puckered three dimensional conformations that allow all the bond angles to be nearly tetrahedral.

UNIVERSITY

#### Course Code : BSCC2001

**Course Name: Organic Chemistry I** 

#### References

- Morrison, R. N. & Boyd, R. N. *Organic Chemistry*, Dorling Kindersley (India) Pvt. Ltd. (Pearson Education).
- 1. Finar, I. L. Organic Chemistry (Volume 1), Dorling Kindersley (India) Pvt. Ltd. (Pearson Education).
- 2. Finar, I. L. Organic Chemistry (Volume 2: Stereochemistry and the Chemistry ofNatural *Products*), Dorling Kindersley (India) Pvt. Ltd. (Pearson Education.
- 3. Eliel, E. L. & Wilen, S. H. Stereochemistry of Organic Compounds, Wiley: London, 1994.
- 4. Kalsi, P. S. Stereochemistry Conformation and Mechanism, New Age International, 2005.
- 5. McMurry, J.E. Fundamentals of Organic Chemistry, 7th Ed. Cengage Learning India Edition, 2013.

**υΝΙνεrsιτ** 

**Course Code : BSCC2001** 

**Course Name: Organic Chemistry I** 

# Thank You

# GALGOTIAS UNIVERSITY

Name of the Faculty: Dr. Diwakar Chauhan