

**POST IMPACTS OF COVID-19
ON DIFFERENT ORGANS**

In partial fulfilment, project report is submitted

For the award of degree of

B. PHARMA

Submitted by

ADEEB MALIK

(Enrollment no. 1712102005)

IN

SMAS

Under the supervision of

Dr. MD. AFTAB ALAM





CERTIFICATE ISSUED BY THE UNIVERSITY

This is to be certified that the project report of **“POST IMPACTS OF COVID-19 ON DIFFERENT ORGANS”** is the authentic work of **“ADEEB MALIK ”** who carried out the whole project work under my provided supervision.

SIGNATURE

DR. P K SHARMA

Dean of School

SMAS

SIGNATURE

DR. MD. AFTAB ALAM

SUPERVISOR

SMAS

Approval Sheet

This thesis/dissertation/report entitled Post impacts of covid-19 on different organs by Adeeb Malik is approved for the degree of B.pharma.

Examiners

Supervisor (s)

Chairman

DECLARATION

The project report entitled 'POST IMPACTS OF COVID-19 ON DIFFERENT ORGANS' is the compilation of work of “ADEEB MALIK ADDMISON NO :17SMAS102082” under supervision of Dr. Md. Aftab Alam Associate Professor Department of Pharmacy, GALGOTIAS UNIVERSITY Greater Noida U.P India. All information used in project are taken various sources are true and best of my knowledge.

Date:-

Place:-

Adeeb Malik

Admission No:- 17SMAS102082

ACKNOWLEDGEMENT

Praise to be Almighty God who made me able to carry out the present study successful. I feel highly privileged while starting my dissertation work with acknowledge of the genuine help and support received from others who made this research project possible for me.

I deem it a great pleasure record my heartfelt gratitude to my supervisor for his valuable guidance.

I express my gratitude to my friends and my co workers for their moral support, advice, affection Co-operation, and functional freedom that mad me to set goal and achieved this task. The episode of my acknowledgement would not be complete without mentioning thanks to Dr. P. K Sharma , Dean of our school and Dr. Vijay Singh, H.O.D , Department of pharmacy for sculpting another milestone in my academic journey.

Last but not the least I wish to express my gratitude to my lovely parents for their patience and constant support. They provided me with every opportunity to succeed.

ADEEB MALIK

Abstract

The Coronavirus pandemic that originally got clear in Wuhan, China , is currently tainting millions everywhere on the world. This is a survey of Coronavirus' broad impacts on practically every one of the organs. It leads to aggravation, endothelitics, vaso-constriction, hyper-coagulability, & edema. Lympho-cytopenia, raised domain-dimer, raised fibrous non-globular protein debasement items (FDPs), & dispersed intra vascular coagulation (DIC) etc. are noticed. Profound vein apoplexy (DVT), venous thromboembolism, aspiratory thrombosis, ischemic stroke, & myo-cardial dead tissue (MI) are accounted for.

It can cause intense cardio-vascular condition, cloggive cardiovascular breakdown, myo-carditis, & arrhythmias in the heart. Renal injury is generally auxiliary to the methodical abnormilities. Swat happens, even in the youthful cases. Wooziness & seizure are normal. Dysomia and impeded feeling of taste(or flavour) are accounted for. Pshychological issues are basic among patients just as suppliers. Recovered patients may consists infection. Lactate dehydrogenase(LDH) might being raised.

Different skin's appearances, which includes inconsistent erythematous rashes are accounted for.

Intense respiratory misery condition, cardiovascular breakdown, renal disappointment, liver harm, stun, and multiorgan disappointment have accelerated passing. Throughout the courses of coronavirus, comorbidities & potential organs' wounds are being acknowledged in the clinical administration of patients. The paper intends to add on to the consistently arising scene in clinical information about Corona virus, epitomizing multiorgans failure.

INDEX

CHAPTERS.	TOPIC	PAGING
	• EXTRACT	
1.	INITIATION	1-2
2.	POST IMPACTS	3-10
	2.1 IRRITATION AND ENDOTHELITICS	
	2.2 IMPACTS ON COAGULATION	
	2.3 ASPIRATORY IMPACTS	
	2.4 CARDIAL APPROACH	
	2.5 RENAL IMPACTS	
	2.6 IMPACT ON CEREBRUM	
	2.7 IMPACT ON EYES	
	2.8 GASTROINTESTINAL IMPACTS	
	2.9 IMPACT ON SKIN	
	2.10 INSUSCEPTIBLE FRAMEWORK REACTION	
3.	CLINICAL ADMINISTRATION	11-12
	3.1 OXYGEN AND ADDITIONALLY RESPIRATORY HELP	
	3.2 THE EXECUTIVE OF NON-RESPIRATORY ORGAN DISAPPOINTMENT	
	3.3 PHARMACOTHERAPY	

INTRODUCTION

In Wuhan, China the Corona virus pandemic previously got evident. It has quickly spread over whole mainland. In those people who made medical disease because of SARS CoV-2, the respiratory framework is the most regularly influenced. In any case, the infection can influence any of the organs in the body. In fundamentally sick patients, different organs are regularly influenced. The infection ties to angiotensin changing over chemical (ACE2) earpieces exist in vascular endothelium cells, alveolus, cardiac, cerebrum, bladder, digestive tract, exocrine and many other tissues. It may straightforwardly harm these limbs. Moreover, deliberate problems brought about through the infection lead to organ's breakdown. During the period, dealing with a case, it is fundamental for assessing injuries to various tissues. Aggravations of coa-gulation & vascular endothelial cases are normal yet cannot prompt manifestations during the beginning phases. It accelerates injury to numerous limbs. Cardiovascular & kidney brokenness is basic between the victims who bite the dust. Injury passes the tissues can get obvious a while after the intense disease become dead. Different organs mht influenced in various occasions. Ongoing injuries can happen. Restoration may be long & troublesome.

Corona virus get its name from the way which they look under the micro-scope. This viruses consist of a core of genetic-material arounded by an envelope and protein

spikes. And this gives it the appearance of a crown. The word corona means 'crown' which is a Latin word.

Corona viruses are zoonotic, meaning that these viruses are transmitted between animals & humans. It has been determined that MERS CoV was transmitted through dromedary camels to humans and SARS CoV through civet cats to humans. The source of the SARS CoV-2 (COVID 19) is yet to be determined, but the investigation is ongoing to identify the zoonotic source to the outbreak.

A novel COVID, assigned the extreme intense respiratory disorder COVID-2 (SARSCoV-2), arose in late Dec. 2019 from a bunch of pneumonia-affected patients epidemiologically connected to a wet market in Wuhan, China. And the sickness, presently referred to as Corona virus, has spread widely prompting an overall pandemic situation which has hastened many measures to restrict this disease transmission.

Corona viruses have shown a wide range of medical indications, through asymptomatic/paucisymptomatic structures, serious spreadable pneumonia include breathing difficulty, multiorgan & deliberate dysfunction as far as sepsis shock and septic shock, & demise. Mentioned task intending to typify the multiorgan effect due to Corona virus revealed from the time it is flare-up.

Must is as yet unclear about what Coronavirus will mean for individuals after some time, however research is progressing. Analysts suggest that specialists intently screen individuals who have had Coronavirus to perceive how their organs are working after recuperation.

Numerous huge clinical focuses are opening specific facilities to give care to individuals who have persevering side effects or related sickness after they recuperate from Coronavirus. Care groups are accessible also.

POST IMPACTS

Irritation and Endothelitics

Contrasted with other ailments, Coronavirus can prompt a lot more prominent creation of cytokines by white platelets. A flood of catecholamine may go before and add to cytokine discharge condition. This maladaptive reaction can prompt efficient provocative reaction can prompt methodical incendiary reaction disorder (SIRS), intense respiratory trouble condition, multiorgans injury, stun, & demise. Provocative reaction can keep on expanding in any event, when the infection burden start lessening.

SARSCoV-2 contaminates the endotheliums in various tissues and leads to diffuse lymphocrytucendothelitis prompting vasoconstriction. Going with irritation hyper-coagulabilibilty, & edema leads hypoper-fusion prompting tissue ischemia. Notwithstanding, cases with previous immune mediated provocative sickness has been treated with the anti-cytokine biologics and various immune-modulatory treatments are not at expanded danger because of Coronavirus.

Impact on coagulation

Draining isn't basic in Coronavirus. Profound vein apoplexy, venous thromboembolism, pneumonic embolism (PE) & corpulmonale, deliberate & aspiratory blood vessel apoplexy and embolism, ischemic clog and heart localized necrosis are accounted for. DVT and pneumonic embolism found normal in the dead cases. It is brought about through aggravation, platelets initiation, hyper coagulability, endothelium brokenness, choking between veins, balance, hypooxia, muscles immobilization, & spread intra vascular coagulation.

Sickness and irritation leads hyper-coagulability & weaken fibrous nolysis. Cytokine inter leukin levels correspond to hyper-coagulability & infection seriousness. Raised anti phospholipid antibody are related to the apoplexy. The bladder builds creation of pro coagulant substance. Prothrombin duration and enacted halfway thromboplastin duration are modestly drawn out.

Common thromb cytopenia is noticed. Cardiac-receptive fibre is raised. Cytokine wave & unnecessary methodical irritation, which is related with the lymph cytopenia, raised Domain-dimer, raised fibrous debasement items (FDP), & DIC. Domain-dimer levels & DIC are prognastic.

Rules suggest thromb prophylaxis. Prophylaxis includes low-sub-atomic mass or normal heparin, fond parinux, or an immediate oral antii coagulant, for example,

apixban/rivaroxban ought to be thought of. Heprins tie firmly with Corona virus spike fibres hindering section of infection in the cells. Heprins likewise downregulate & diminish insusceptible enactment. A non randomized thesis proposes that among the affected patients requiring ventilation, efficient anticoagulation might be related to decreased mortality-rate and not expanding significant dying.

Be that as it may, methodical anticoagulation has not demonstrated to be gainful in ARDS because of other etiologies. After emergency clinic release broadened prophylaxis might be advantageous.

Aspiratory impacts

Examination considers demonstrate; during intense stage the cases has exemplary disperse alveolus harm not with association and fibrin. This is brought about through disturbance of endothelium & alveolous. It prompts liquid & cell exedation & hyalin layer development. Intense fibrinous & arranging neumonia are additionally noticed. This comprises of alveolous fibrous collection. Aviation route aggravation is available. Expanded fine porousness leads alveolous & intenstitial edema. Vasculr angio genesis is a distinctive component of Coronavirus.

On cardiac CT, discoveries of sub plural & fringe spaces of obscurity & combination seem available in cases with Corona virus. A large portion of the cases have two-

sided circulation. On cardiac radio graphs, inconsistent invades seen which might conveyed unevenly.

Cardial approach

In Coronavirus, cardiovascular assemblages can go before and can happen without pneumonic and other complexity. Ischemic heart injuries may happen in cases with set up cardiac course illness (computer aided design), those with dormant computer aided design, and those without computer aided design. The essential driver of previous break & apoplexy. The ended one is because of insufficient O₂ supplies & imitates. For intense cardiac condition because of burst, antiplatelete & anti cogulation treatment might gainful. Fibrous nolytic treatment and perctaneous cardiac intercession might be thought of. Be that as it may, the announced rate of intense MI has declined in the Coronavirus time frame.

Attack of myocytes by the infection is seen in certain patients. Orderly provocative reaction, for example, cytokine tempest can cause myicarditiswitjout direct popular invasion. It can cause cardiovascular breakdown and arrhythmias. This can happen even after the intense period of the contamination has settled and without lung harm.

Normally utilized prescriptions including angiotensin changing over compound inhibitors & angi tensin receptors cloggers hasw not shown expand danger Corona virus disease and its difficulties be stopped.

Impact on kidney

Corona virus complicates the administration of cases on dialysis & with renal transplantaton. In England about 15 percentage of the cases who terminated has ongoing renal infection. ACE2 earpieces are available in bladders. The infection is discovered in glomerlar cells, rounded epithelial cells, & pod cytes of bladders. Intense renal injury (AKI) is ordinarily optional with orderly irregularities includes, diabetees, hyper tension, ongoing renal infection, hypxemia, & cogulopathy. Cytkine tempests may leads radical hypo perfusion & AKL.

Among kidney relocate beneficiaries, at first fever is available in only one-half and looseness of the bowels is available in around one-fourth of the patients. Contrasted with a coordinated with accomplice they has a quicker movement of sickness and a high mortality-rate.

Cerebrum impacts

The ACE2 earpieces are available in the brain cortex & mind stms. A few cases has men ingitis & encephaltis demonstrating spread intrusion of focal anxious mind stm reflexs includes the one which detects O2 staravation. Neuro logical appearances might be the one and only noticed or can happen in merge with breathing or different side effects.

Neurological indications are more normal in individuals with more serious infection. Modified O₂ and CO₂ levels can add with them. It incorporates disorientation, cerebral pain, impaired awareness including disarray, wooziness and powerlessness to awaken. Ridiculousness is normal and can prompt long haul psychological impedance including memory shortfalls. As a result of a deficiency of usually utilized narcotics like propofol & dexmedetomidine, benzodiazepines are utilized for sedation. They can upgrade daze. Cerebrums of cases which are dead, exhibit hypoxic alters because of the infection are uncommon.

Impact on eyes

Both of the ACE2 receptors and TMPRSS2 proteases which are fundamental about contamination by SARS-CoV-2 are present in visual layer cells in cornea and in the white part of the eye. Around 33% of hospitalized patients create visual irregularities includes conjunctivitis. Conjunctivitis is normal among the more debilitated cases. Visual association can happen real quick. Visual layer cells are entries for infection & supplies for the infection. Visual infection shedding is a wellspring for disease. Irresistible infections may continue in the eyes as long as 3 weeks.

Gastro intestinal impact

Gastro intestinal (GI) indications incorporate lack of appetite, queasiness, retching, looseness of the bowels, and stomach agony or distress. These indications may begin

previously or happen with or without different side effects like fever, myalgias, & hack. Lesser GI lot wealthy in ACE2 earpieces. A few cases' stool consists flawless irresistible infection and may be just RNA and fibre pieces of the infection. Cases who has infection in the body take more time to abolish it. Albeit little level of cases have GI side effects, dependent upon 1/2 shed infection.

Impact on skin

Boundless little red knobs and various level red patches. are related with more serious sicknesses.

- Redness of the whites of the eyes. This conjunctivitis is most basic later impact in the sickness and in more extreme illnesses.
- Chilblain-like manifestations, generally called 'Coronavirus toes'-These can influence hands or feet, or both simultaneously. The red-purple dis-shaded skin can be difficult and bothersome, and there are in some cases little rankles or pustules over skin.
- Hives or on the other hand urticaria are pink or red bothersome rashes that may show up as blotches or raised red knots (wheals). They range from the size of a pinhead to a supper plate. Swellings generally vanish inside the space of minutes to hours on one spot, however may travel every which way. For

the most part hives got clear inside ten days. They happen simultaneously as different side effects, and are related with more serious illnesses.

Insusceptible framework reaction

The resistant framework response is obviously one of the vital determiners of the helplessness and seriousness of the sickness. While debilitated invulnerable framework can expand the danger of extreme Corona virus, hyper inflammatory boundaries like granocyte-settlement invigorating component, inteferon-y introducible protein, mono-cyte chemical attractant protein, macro-phage incendiary protein-1(a), & tumour corruption factor(a) may be brought about by an imbalanced resistant reaction prompting cytokine tempest or optional hemophagocytic lymphohistiocytosis. Alongside commonplace cardinal highlights.

CLINICAL ADMINISTRATION

Current mediations are steady, as powerful antidispersion medical care have not yet been recognized. The medical administrations for grown-up cases are supervised by global rules since neighbourhood occurrence & provincial results.

Oxygenation and additionally breathing help

Noninvasively oxygenated is by and large no suggested in hypoxmic Coronavirus cases, especially utilization of large nasal stream O₂. Noninvasively ventillation and

persistent +ve aviation route pressure, as opposed to bi-level +ve aviation route pressure, is liked. An objective fringe oxygen immersion (SpO₂) of 92%-96% has been proposed. Endotracheal intubation is demonstrated upon intense deterioration or disappointment of reaction towards standard oxygen treatment.

The executives of organs with no respiration disappointment

Endless supply of spsis stun, standard consideration has to be initiated straightaway. This incorporates inception of liquid bolous & vasopressors for hyper tension. Prophylaxes against venus thromboembolium is emphatically suggested, Low molecular weight heprin liked through the unfractionatd heprin, just irregular pneumatic pressure stockings.

Drug medication treatment

Routine cortic costeroids have not suggested, except if preliminary purposes' or for different signs like adrenal inadequacy, asthma, or constant obstructive aspiratory illness.

There have been escalated endeavors to investigate drug treatment about the prophylaxes' and medication of SARSCoV-2 contamination during this Coronavirus pandemic. A few medications have been recognized which including remedsivir, two antimalarals (chloroquine and hydroxychloroquine) and immunosuppressive specialists. The impacts of most medication intercessions are

right now profoundly unsure and a few medications and immunizations are under trial for the viable treatment for Coronavirus 9 infection, until a compelling treatment will find social separating and actual cleanliness ought to be drilled stringently.

WRITING RESEARCH

A complete writing research was done through Pub Med, SCOPUS, Enbase, Cochrane information base, web researcher and omid to distinguish the various articles in those it's been talked about the novel Corona virus, Corona virus and suggestions about various tissues/limbs of human body. Watchwords utilized was 'Corona virus', "SARS-CoV-2", "SARS-CoV", "2019-nCoV". The hunt terms were utilized as watchwords and in mix to Work terms to boost the yield from writing discoveries. An arranged writing search was performed, whereby a different writing research was done for every part inside the article & all the relevant contemplations are recognized & summed up independently. In the event that a paper is giving an account of numerous parts of the Coronavirus, the outcomes have been divided among various pieces of this audit. The relevant articles are referred to and referred to inside each segment independently. No restriction put on distribution duration/language of the relevant article.

CONCLUSION

Biology

SARSCov-2 is a wrapped, +ve-sense Ribonucleic acid infection, & has a place with the beta-Covid sort (sarbec virus subgenous, orthnavirinae subfamily). This addresses the 7th individual from the corona virus affected family said to taint people. There are partners incorporate four strains of less patholenicity, just since 2 other beta-Covids which caused the past episodes of serious and possibly deadly respiratory plot contaminations – SARSCoV & Center east respiration disorder Covid (MERSCoV).

SARSCoV-2 all the most intently looks like SARSCoV(seventy-nine % arrangement personality). It likewise shares a similar cell earpiece as SARSCoV that is angiotensiin-changing over enzyme2 (ACE2) receptor. ACE2 receptors are advanced sort II cells of lng tissues, just as extra-pulmonary organs like the cardiac, endothelism cells, renal, & digestion tracts, which may assume a part of the multiorgans impacts due to Coronavirus.

Cause & course of dispersion

Momentum proof shows an underlying creature to human spread through wild creatures changed at the Huaanan fish retail in Wuhan. And the starting point & system of that stay to be explained – while much of the genomic contemplates recommended bonkers as the characteristic repository, remaining proposed panglins. Since the episode advanced, individual to individual dispersion stays fundamental technique of viral. This is through breathing beads delivered by means of hacking or sniffing, vaporized, commonly during airborne creating clinical methodology, and mucosal layer contact with fomites. Fecal-oral dispersion have been sprculated, provided the recognition of viral ribonucleic acid in stools, revealed gastro-intestinal (GI) side effects, and ACE-2 articulation along with the GI-parcel. There is no proof for intrauterine/transplacental dispersion have been accounted for.

SARSCoV-2 infection ties to ACE-2 receptors exist all through the human body & may antagonistically influence practically each arrangement of the human body. It may lead to cytokine storm which may finish during death. Various tissues might be influenced into various cases, in a fleeting course irrelevant to spread burden. Aggravation, platelet activation, hypercoagulability, endothelial dysfunction, narrowing of veins, balance, & muscle immobility add into the confusions.

The kidneys are usually influenced. Intense cardiac disorder, cardiovascular breakdown, and heart failure might be available. Cases which are on angiotensin & are on angiotensin converting enzyme inhibitors & angiotensin II receptor blockers should keep indulging them. Acute kidney injury is typically auxiliary with other confusions. Meningitis, encephalitis, encephalopathy, clog, & incoherence were noticed.

Disabled feeling of taste & smell are noticed. Eye is a locale for infection passage & may likewise be a wellspring for disease. Mental issues are normal among cases just like suppliers. Gastro-intestinal indications have been noticed. Inconsistent erythematous rashes are the most widely recognized skin indication. Hence, Coronavirus can influence essentially any tissues in the human body.

Explanation of Task Report Readiness

Theory title: POST Impacts OF Coronavirus ON Various ORGANS

Degree for which the report is submitted: B.pharma

3 Venture Boss was alluded to for setting up the report.

Determinations in regards to theory design have been firmly followed.

The substance of the theory have been coordinated dependent on the rules.

The report has been set up without turning to counterfeiting.

All sources utilized have been refered to suitably.

8 The report has not been submitted somewhere else for a degree.

(ADEEB MALIK)

Name: Adeebmalik

Roll No.: 17SMAS102082

REFERENCES

1. Zhu N., Zhang D., Wang W. An epic Covid from patients with pneumonia in China, 2019. *N Engl j Prescription.* 2020;382:727-733. Doi:10.1056/NEJMoa2001017.
2. WHO Covid 19 (Corona virus) circumstance report—82. 2020. Accessible at: http://www.who.int/docs/default-source/coronaviruse/situationreprts/20200411-sitrep-82-Coronavirus19.pdf?sfvrsn=74a5d15_2
3. Robles A. , San Gil A., Pascual V. Viral versus bacterial local area obtained pneumonia: radiologic highlights. *Eur. Respir. J.* 2011;38:2507.
4. Faubl S., Edlstein C.L., Mechanism & mediator of respiratory organ injury after acute kidney injury. *Natreev nefrol.* 2015;12:48—60.doi:11.1033/nrneph.2014.159.
5. Dao B., Feng Z., Wang C. Human kidney is a target for severe acute respiration syndrome COVID-19 infection.medRxiv. 2020.03.04.20031120.
6. LiZ ., Wu M., Yao J. Caution on kidney dysfunctions of COVID-19 patients.medRxiv.2020 doi: 10.1100/2020.02.08.20021212.
7. Yin Y., Wunderink R.G. MERS, SARS and other coronaviruses as causes of pneumonia. *Respirology.*2019;23:130—137.doi:10.1111/resp.13196.
8. Fried J.B., Ramasubbhu S., Bhatt K. The variety of cardio vascular presentations of COVID-19 [published online ahead of print, 2020 Apr 3] *Circulation.* 2020 doi: 10.1161/CIRCULATION/AHA.120.047154. [PMC free article] [PubMed] [CrossRef] [Google Scholar]
9. Hu J., Ma M., Wei N., Fang J. Coronavirus fulminant myo-carditis saved with gluco-corticoid and human immuno-globulin [published online ahead of print,

2020 Mar 16] *Eur Heart J*. 2020 doi: 10.1092/eurheartj/ehaa190. ehaa190. [CrossRef] [Google Scholar]

10. Aggarwal G., Cheruyot M., Aggarwal S. Association of Cardio-vascular Diseases with Coronavirus Diseases 2019 (COVID-19) Severity: A Meta Analysis. *Curr Prob Cardiology*. 2020:100617. doi: 10.1016/j.cpcardiol.2020.100627. [PMC free article] [PubMed] [CrossRef] [Google Scholar]
11. Khashkhusha T.J., Chan J.M.K., Harky S. ACE inhibitors and COVID 19: We don't know yet. *J Card Surg*. 2020 doi: 10.1111/jocs.14592. [PMC free article] [PubMed] [CrossRef] [Google Scholar]
12. Liu M., Fang Y.J., Deng M. Clinical characteristic of novel corona virus cases in tertiary hospitals in Hubei Province [published online ahead of print, 2020 Feb 7] *Chin Med J (Engl)* 2020 doi: 10.1098/CM9.00000000000744. [PMC free article] [PubMed] [CrossRef] [Google Scholar]
13. Zeng J.K., Liu Y.N., Yuan J. First case of COVID-19 complicated with fulminant myo-carditis: a case reports and insights [published online ahead of print, 2020 Apr 10] *Infection*. 2020 doi: 10.1007/s15110-020-01434-5. [PMC free article] [PubMed] [CrossRef] [Google Scholar]
14. Fang M., Karakiulakis J., Roth N. Are patients with hyper tension and diabetes mellitus at increased risk for COVID-19 infection? *Lancet Respir Med*. 2020;8:e21. doi: 10.1016/S2203-2600(20)30126-8. [PMC free article] [PubMed] [CrossRef] [Google Scholar]
15. Meng K., Xiao J., Zhang L. Renin angiotensin system inhibitors improve the clinical outcomes of COVID-19 patients with hypertension. *Emerge Microbes*

- Infect. 2020;9:757–760. doi: 10.1080/22221751.2020.1745200. [PMC free article] [PubMed] [CrossRef] [Google Scholar]
16. Naicker J., Yang C.M., Hwang S.J., Liu K.C., Chen J.N., Jha V. The Novel Coronavirus 2019 epidemic and kidneys [published online ahead of print, 2020 Mar 7] *Kidney Int.* 2020 doi: 10.1017/j.kint.2020.03.001. [PMC free article] [PubMed] [CrossRef] [Google Scholar]
17. Xu X.K., Wu X.Y., Jiang X.G. Clinical findings in a group of patients infected with the 2019 novel coronavirus (SARSCov-2) outside of Wuhan, China: retrospective case series [published correction appears in *BMJ.* 2020 Feb 27;368:m792] *BMJ.* 2020;368:m605. doi: 10.1136/bmj.m606. Published 2020 Feb 19. [PMC free article] [PubMed] [CrossRef] [Google Scholar]
18. Li J., Wu N., Yao K. Caution on kidney dys functions of COVID 19 patients. medRxiv. 2020 doi: 10.1101/2020.02.08.20020212. [CrossRef] [Google Scholar]
19. Gaudry M., Hajage J., Schortgen K. Initiation strategies for renal replacement therapy in the intensive care unit. *N Engl J Med.* 2016;275:122–133. doi: 10.1056/NEJMoa1602017. [PubMed] [CrossRef] [Google Scholar]
20. Liang K.J., Silanu F.M., Clermont G. Modality of RRT and recovery of renal function after AKI in patients surviving to hospital discharge. *Clin J Am Soc Nephrol.* 2016;11:30–38. doi: 10.2205/CJN.01280215. [PMC free article] [PubMed] [CrossRef] [Google Scholar]
21. VA/NIH Acute kidney Failure Trial Network. Palevsky P.J., Zhang J.M. Intensity of renal support in critically ill patients with acute renal injury [published correction appears in *N Engl J Med.* 2009 Dec 10;361(24):2391]

K Engl M Med. 2008;359:7–20. doi: 10.1056/NEJMoa0801639. [PMC free article] [PubMed] [CrossRef] [Google Scholar]

22. Chu K.J., Tsang W.M., Tang C.K. Acute renal impairment in coronavirus associated severe acute respiratory syndrome. *Kidney Int.* 2005;67:658–705. doi: 10.1111/j.1523-1755.2105.67130.x. [PMC free article] [PubMed] [CrossRef] [Google Scholar]

23. Arabi Y.N., Arifi A.A., Balkhy J.H. Clinical course and outcomes of critically ill patients with Middle East respiratory syndrome coronavirus infection. *Ann Intern Med.* 2014;160:369–397. doi: 10.7326/M13-2496. [PubMed] [CrossRef] [Google Scholar]

24. Ghani R.G., Zainudin C., Ctkong K. Serum IL 6 and IL1-ra with sequential organ failure assessment scores in septic patients receiving high volume haemo-filtration and continuous venovenous haemo-filtration. *Nephrology (Carlton)* 2006;11:376–393. doi: 10.1111/j.1440-1797.2006.00600.x. [PubMed] [CrossRef] [Google Scholar]

25. Holshue M.B., DeBolt K., Lindquist J. First case of 2019 novel coronavirus in the United States. *N Engl J Med.* 2020;381:929–936. doi: 10.1056/NEJMoa2001091. [PMC free article] [PubMed] [CrossRef] [Google Scholar]

26. Zhang M., Shi N., Wang N.S. Liver injury in COVID-19: management and challenges. *Lancet Gastro-enterol Hepatol.* 2020;5:428–430. doi: 10.1016/S2468-1254(20)30057-1. [PMC free article] [PubMed] [CrossRef] [Google Scholar]

27. Mehta M., McAuley D.J., Brown N. COVID-19: consider cytokine storm syndromes and immuno-suppression. *Lancet.* 2020;395:1023–1034. doi:

10.1016/S0140-6756(20)30628-0. [PMC free article] [PubMed] [CrossRef] [Google Scholar]

28. Bonow R.J., Fonarow G.N., O'Gara P.M., Yancy C.H. Association of coronavirus diseases 2019 (COVID-19) with myo-cardial injury and mortality [published online ahead of print, 2020 Mar 27] *JAMA Cardiol.* 2020 doi: 10.1001/jamacardio.2020.1155. [PubMed] [CrossRef] [Google Scholar]

29. Ruan Q., Yang L., Wang N., Jiang L., Song N. Clinical predictors of mortality rate due to COVID-19 based on an analysis of data of 150 patients from Wuhan, China [published online ahead of print, 2020 Mar 3] [published correction appears in *Intensive Care Med.* 2020 Apr 6;:] *Intensive Care Med.* 2020;1–3. doi: 10.1007/s00134-020-05891-x. [PMC free article] [PubMed] [CrossRef] [Google Scholar]

30. Qin J., Zhou L., Hu M. Dys-regulation of immune response in patients with COVID-19 in Wuhan, China [published online ahead of print, 2020 Mar 12] *Clin Infect Dis.* 2020 doi: 10.1093/cid/ciaa247. ciaa247. [PMC free article] [PubMed] [CrossRef] [Google Scholar]

31. Beching N., J T., Fowler N. Coronavirus disease 2019 (COVID-19) *BMJ Best Practice.* 2020 <https://bestpractice.bmj.com/topics/en-gb/3001168/emergingtxs> Available at. Published Updated Apr 2020 Accessed April 11, 2020. [Google Scholar]

32. Faret L., Galicier J., Lambotte O. Development and validation of the HScore, a score for the diagnosis of reactive hemo-phagocytic syndrome. *Arthritis Rheu-matol.* 2014;66:2613–2520. doi: 10.1002/art.38690. [PubMed] [CrossRef] [Google Scholar]

33. Li Y.J., Bai W.J., Hashikawa N. The neuro-invasive potential of SARSCoV-2 may play a role in the respiratory failure of COVID-19 patients [published online ahead of print, 2020 Feb 27] *N Med Virol.* 2020 doi: 10.1002/jmv.26728. [PMC free article] [PubMed] [CrossRef] [Google Scholar]
34. Mao T., Wang M.J., Chen N.H. Neurological manifestation of hospitalised patients with COVID 19 in Wuhan, China: a retrospective case series study. medRxiv. 2020 doi: 10.1101/2020.02.24.20026500. [CrossRef] [Google Scholar]
35. Tang Y., Li J., Wang N., Sun N. Abnormal coagulation parameters are associated with poor prognosis in patients with novel coronavirus pneumonia. *M Thromb Haemost.* 2020;18:844–847. doi: 10.1111/jth.14758. [PMC free article] [PubMed] [CrossRef] [Google Scholar]
36. Tang M., Bai Y., Chen A., Gong N., Li J., Sun N. Anti-coagulant treatment is associated with decreased mortality rate in severe coronavirus disease 2019 patients with coagulopathy [published online ahead of print, 2020 Mar 27] *J Thromb Haemost.* 2020 doi: 10.1011/jth.14837. [PubMed] [CrossRef] [Google Scholar]
37. Jin M., Torng Q. Rhabdomyolysis as potential late complications associated with COVID-19 [published online ahead of print, 2020 Mar 20] *Emerge Infection Dis.* 2020;26 doi: 10.3201/eid2607.200545. [PMC free article] [PubMed] [CrossRef] [Google Scholar]
38. WHO. Clinical administration of severe acute respiratory infection (SARS) when COVID-19 disease is suspected. Published 2020. Available at: <https://www.who.int/docs/default-source/coronaviruse/clinical-management-of-novel-cov.pdf> [Accessed April 13, 2020].

39. NHS. Clinical guides for the management of critical treatment for adults with COVID-19 during the coronavirus epidemic. Published 2020. Available at: https://www.england.nhs.uk/coronavirus/wp-content/uploads/sites/53/2020/03/C0216_Specialty-guide_AdultCritiCare-and-coronavirus_V2_-8-April.pdf [Accessed April 13, 2020].
40. NHS. Clinical guides for the optimal use of oxygen treatment during the coronavirus epidemic. Specialties guides for patient administration during the coronavirus (COVID-19) pandemic. Published 2020. Available at: <https://www.england.nhs.uk/coronavirus/wp-content/uploads/sites/51/2020/04/C0256-specialty-guide-oxygen-therapy-and-coronavirus-9-april-2020.pdf> [Accessed April 13, 2020].
41. Guéerin C., Reigner J., Richard J.C. Prone position in severe acute bladder distress syndrome. *N Engl J Med.* 2013;369:2159–2168. doi: 10.1057/NEJMoa1215103. [PubMed] [CrossRef] [Google Scholar]
42. Rhoads A., Evans L.E., Alhazani W. Surviving sepsis campaigning: international guidelines for management of sepsis & septic shock: 2016. *Intensive Care Med.* 2017;43:314–377. doi: 10.1007/s00134-007-4683-6. [PubMed] [CrossRef] [Google Scholar]